

CHILDREN'S EXPOSURE TO RARE EARTH ELEMENTS AND CANCER RISK

SADEG N. ^{1,2}, BROUSSE F. ¹, MALONGA G. ¹, POMMIER A. ^{2,4} and BELHADJ-TAHAR H. ^{2,3}

¹ International Medical Institute of Toxicology and Pharmacology, rue Henri Becquerel, 60230 Chambly, France - ²toxSeek Scientific Board, 95300 Ennery, France
³ Research and Expertise group, French Association for Medical Research Advancement (AFPREMED), 31100 Toulouse, France - ⁴ MSA Picardie, 31100 Toulouse, France

BACKGROUND

Rare earth elements (REE) are extensively used in the renewable energy technologies such as **wind turbines, batteries, catalysts** and **electric cars**.

While there are many studies about cancer epidemiology of several heavy metal, much less is known about rare earth elements despite the fact that the lanthanides included in rare-earth metals have shown **interactions with cellular metabolism and physiology** due particularly to their catalytic properties (impact on phosphokinase) and their radii ionic similitude with calcium (impact on Calmodulin) ^[1].

Our team have recently developed ICP-MS technics for hair metallic trace element screening applied to epidemiology investigation on pediatric cancer ^[2].

In this abstract, we describe the first investigation on hair REE concentration in pediatric cancer among children of the Sainte-Pazanne in northern France (fig. 1).



(fig. 1)

METHODS


Subjects: 64 children aged 5 to 15 years were included in this investigation; 20 from Sainte-Pazanne (6 500 habitants) in the Loire-Atlantique department in western France and 44 healthy children from all other part of France.
Among 20 children of Sainte-Pazanne 12 suffered from cancer.

Briefly, 10-50 mg of hair was washed twice and hydrolyzed in nitric acid medium during 12h. The diluted hydrolysates and External standards were tested by ICP/MS Injection Agilent 7800. Rare Earth element dosed: Lanthane (La), Cerium (Ce), Praseodyme (Pr), Neodyme (Nd), Samarium (Sm), Europium (Eu), Gadolinium (Gd), Dysprosium (Dy), Holmium (Ho), Erbium (Er), Thulium (Tm) and Ytterbium (Yb).

Results: 4 rare earth elements (Pr, Dy, Ho and Tm) were found significantly in large proportion in Sainte-Pazanne children compared to the control population: 231% Pr, 257% Dy, 327% Ho and 400% Tm respectively (n = 64, p <0.05) (Fig 2).

Within the population of the target area, the Praseodyme (Pr), and Holmium (Ho) levels were significantly high in 20% (4/20) of sick children: Pr (Patient: 0.033 ±0.01 µg/g versus healthy: 0.001 ± 0.001 µg/g) and Ho (Patient: 0.003 ±0.002 µg/g versus healthy: 0.001 ± 0.001 µg/g).

Conclusion: This results suggests that the prevalence of children cancer in lanthanide contaminated area might be much higher than elsewhere and rare earth elements could contribute to pathogenesis of pediatric cancer.



(fig. 2 : Comparative hair REE concentration in Sainte-Pazanne's subjects versus Control group)

Statement: The authors declare that there is no conflict of interest.

References:

- EVANS, C.H., 1990. Biochemistry of the lanthanides. Plenum Press, New York.
- BROUSSE and Coll. Assessment of Exposure to Organic and Inorganic Pollutants in Children's Hair International Journal of Public Health Research 2019; 7(1): 18-22.